

Chronic Phase Chronic Myelogenous Leukemia (CP-CML) Pathways

Patient Name: _____ Date of Birth: _____
Member Number: _____ Treatment Start Date: _____
Pathology: _____ Stage: _____
Line of Therapy: _____ ICD-10 Code: _____

Biomarkers/Characteristics: (select all that apply)

CML Phase: ___ Chronic Phase ___ Accelerated Phase ___ Lymphoid Blast Phase ___ Myeloid Blast Phase ___ Not Reported

Imatinib resistant or intolerant: ___ Yes ___ No Philadelphia chromosome: ___ Positive ___ Negative

T315I: ___ Positive ___ Negative

Mutation: ___ V299L ___ T315I

New Diagnosis of CML

- Low, Intermediate, or High-Risk Disease*
 - ☐ Imatinib (Gleevec)
- Intermediate or High-Risk Disease*
 - ☐ Dasatinib (Sprycel)
 - ☐ Nilotinib (Tasigna)

Second Line of Therapy (2nd Line)

- Resistant disease to primary treatment, Suboptimal Response[†], or Intolerance to 1st Line
 - ☐ Bosutinib (Bosulif)
 - ☐ Dasatinib (Sprycel)
 - ☐ Nilotinib (Tasigna)
- Presence of T315I mutation
 - ☐ Ponatinib (Iclusig)

* For patients with intermediate or high risk disease based on Sokal or Hasford score:

Sokal: Intermediate Risk=0.8-1.2; High Risk>1.2

Hasford: Intermediate Risk=781-1480; High Risk>1480

[†] Defined as lack of complete hematologic response or BCR-ABL1 transcripts > 10% (IS) or lack of partial cytogenetic response on bone marrow cytogenetics.

Note: Pathways are independent of specific health plan medical policy coverage criteria. Health plan medical policy/clinical guidelines should be consulted to determine whether proposed services will be covered. Biosimilars of reference products listed are considered "on pathway." However, reimbursement for biosimilar products may be impacted by health plan specific formularies, medical policy and preferred product rules.