

Testicular (Germ Cell Tumors) Cancer Pathways

Patient Name: _____

Date of Birth: _____

Member Number: _____

Treatment Start Date: _____

Pathology: _____

Stage: _____

Line of Therapy: _____

ICD-10 Code: _____

Adjuvant Therapy

- Stages II-IIIC and IS
 - **Seminoma**
 - BEP:** bleomycin, etoposide, and cisplatin
 - EP:** etoposide and cisplatin
 - **Non-Seminoma** after Retroperitoneal Lymph Node Dissection (RPLND)
 - EP:** etoposide and cisplatin

First Line of Therapy (1st Line)

- Stages II-IIIA and IS
 - **Seminoma and Non-Seminoma**
 - BEP:** bleomycin, etoposide, and cisplatin
 - EP:** etoposide and cisplatin
- Stages IIIB and IIIC
 - **Seminoma** with Good and Intermediate Risk*
 - BEP:** bleomycin, etoposide, and cisplatin
 - **Seminoma** with Good Risk
 - EP:** etoposide and cisplatin
 - **Non-Seminoma**
 - BEP:** bleomycin, etoposide, and cisplatin

* BEP is typically given for 3 cycles in good risk seminoma, and 4 cycles in intermediate risk.

† EP given for 4 cycles

Note: Pathways are independent of specific health plan medical policy coverage criteria. Health plan medical policy/clinical guidelines should be consulted to determine whether proposed services will be covered. Biosimilars of reference products listed are considered "on pathway." However, reimbursement for biosimilar products may be impacted by health plan specific formularies, medical policy and preferred product rules.



8600 West Bryn Mawr Avenue
South Tower – Suite 800 Chicago, IL 60631

Last review: 5/16/2023 | Effective date: 5/16/2023

© 2023 Carelon Medical Benefits Management