Gastric, Esophageal, and Gastroesophageal Junction Adenocarcinoma Pathways

Patient Name:	Date of Birth:
Member Number:	Treatment Start Date:
Pathology:	Stage:
Line of Therapy:	ICD-10 Code:
Will the patient be undergoing surgery?YesNo	Will the patient be undergoing radiation therapy?YesNo
Neoadjuvant Therapy (Pre-Operative, Peri-Operative, Primary Therapy)	
 Potentially Resectable Disease 	
☐ Cisplatin and fluorouracil (5FU)	
☐ Fluorouracil (5FU) and cisplatin with concurrent radiation therapy (RT)	
☐ FLOT: Fluorouracil (5FU), leucovorin, oxaliplatin, and docetaxel (Taxotere)	
☐ Paclitaxel and carboplatin with concurrent RT*	
Adjuvant Therapy (Post Operative-Adjuvant) – Resected Disease	
☐ Fluorouracil (5FU) and leucovorin with concurrent RT	
☐ Nivolumab (Opdivo)*	
First Line of Therapy	
 Locally Advanced, Metastatic, or Recurrent Diseas 	se
 Unresectable, HER2 Positive 	
☐ Cisplatin, fluorouracil (5FU), and trastuz	zumab
 HER2 Negative 	
☐ Fluorouracil (5FU) and Cisplatin [†]	
☐ Fluorouracil (5FU) +/- Leucovorin and irinotecan (Camptosar) FOLFIRI	
☐ FLO/FOLFOX: fluorouracil (5FU), leucovorin, and oxaliplatin	
☐ FLP: fluorouracil (5FU), leucovorin, and cisplatin	
□ FOLFOX + nivolumab: fluorouracil (5FU), leucovorin, oxaliplatin, and nivolumab (Opdivo) (CPS ≥ 5)	
□ Pembrolizumab (Keytruda), fluorouracil (5FU) and cisplatin (CPS ≥10)	
□ Pembrolizumab (Keytruda), capecitabine (Xeloda) and oxaliplatin (CPS ≥10)	
Second Line of Therapy (2nd Line)	
Unresectable Locally Advanced, Metastatic, or Recurrent Disease	
☐ Irinotecan (Camptosar)	
□ Paclitaxel	
☐ Trastuzumab deruxtecan (Enhertu)‡ – ((HER2 Positive Only)

Note: Pathways are independent of specific health plan medical policy coverage criteria. Health plan medical policy/clinical guidelines should be consulted to determine whether proposed services will be covered. Biosimilars of reference products listed are considered "on pathway." However, reimbursement for biosimilar products may be impacted by health plan specific formularies, medical policy and preferred product rules.



^{*} Limited to esophageal and gastroesophageal junction cancers only-maximun duration of treatment is one year.

[†] Limited to gastric tumors only

[‡] Use only after prior trastuzumab based therapy