Ovarian Cancer (Epithelial) Pathways

Patient Name:	Date of Birth:
Member Number:	Treatment Start Date:
Pathology:	
Line of Therapy:	
Biomarkers/Characteristics: (Select all that apply) Platinum Sensitive [*] :YesNoNot Reported	Paclitaxel-resistant:YesNo Platinum Refractory/Resistant:YesNoNot Reported
Adjuvant Therapy	
• Stages IA/IB (Grade 2 or 3), and IC (Grade 1-3	3)
Carboplatin and paclitaxel	
Adjuvant, Neoadjuvant, or Primary Therapy	
Stages II-IV	
Carboplatin and paclitaxel (Admini	stered weekly or every 3 weeks)
Metastatic / Recurrent Disease	
Platinum Sensitive*	
□ Carboplatin	
Carboplatin and gemcitabine (Gem	zar)
Carboplatin and paclitaxel	
Carboplatin and weekly paclitaxel	
Platinum Resistant	
Bevacizumab monotherapy	
Docetaxel (Taxotere)	
Gemcitabine (Gemzar)	
Liposomal doxorubicin (Doxil)	
Paclitaxel (weekly)	
Paclitaxel and bevacizumab	
Maintenance Therapy	
After response to initial treatment for platinum-	sensitive disease
 Sensitive* to platinum-based therapies 	
Niraparib (Zejula)	
Olaparib (Lynparza)	
Rucaparib (Rubraca)	

* Platinum sensitive disease is defined as recurrence of greater than 6 months after prior platinum-based therapy

Note: Pathways are independent of specific health plan medical policy coverage criteria. Health plan medical policy/clinical guidelines should be consulted to determine whether proposed services will be covered. Biosimilars of reference products listed are considered "on pathway." However, reimbursement for biosimilar products may be impacted by health plan specific formularies, medical policy and preferred product rules.

