

Lung Cancer: Non-Small Cell Lung Cancer (NSCLC) Pathways

Patient Name: _____

Date of Birth: _____

Member Number: _____

Treatment Start Date: _____

Pathology: _____

Stage: _____

Line of Therapy: _____

ICD-10 Code: _____

Biomarkers/Characteristics: (Select all that apply)

PD-L1 expression: __ Less than 50% __ Equal to or greater than 50%

Actionable Oncogenic Targets*: __Yes __No

Chemoradiation for Localized Disease

- Stages IA-IIIc – Definitive Concurrent Chemoradiation
 - Cisplatin and etoposide
 - Paclitaxel and carboplatin

Adjuvant Therapy

- Stages IB-IIIb
 - Carboplatin and paclitaxel
 - Cisplatin and gemcitabine (Gemzar)
 - Cisplatin and pemetrexed (Alimta)
 - Cisplatin and vinorelbine (Navelbine)

First Line of Therapy (1st Line) – Stages IIIB-IV, and Recurrent

- Squamous and Non-Squamous Cell Carcinoma
 - PD-L1 Expression (TPS) greater or equal to 50%, **without known actionable oncogenic targets***
 - Cemiplimab-rwlc (Libtayo)
 - Pembrolizumab (Keytruda)†
 - Ineligible for Immunotherapy
 - Carboplatin or cisplatin and paclitaxel
 - Carboplatin or cisplatin and gemcitabine (Gemzar)
 - ALK Rearrangement Positive
 - Alectinib (Alecensa)
 - Lorlatinib (Lobrena)
 - EGFR exon 19 deletion or exon 21 L858R mutation positive
 - Osimertinib (Tagrisso)
 - Carboplatin or cisplatin, pemetrexed, and osimertinib (Tagrisso) **(Non-Squamous Only)**

Note: Pathways are independent of specific health plan medical policy coverage criteria. Health plan medical policy/clinical guidelines should be consulted to determine whether proposed services will be covered. Biosimilars of reference products listed are considered “on pathway.” However, reimbursement for biosimilar products may be impacted by health plan specific formularies, medical policy and preferred product rules.

First Line of Therapy (1st Line) – Stages IIIB-IV, and Recurrent - Continued

- Non-Squamous Cell Carcinoma Only
 - PD-L1 Expression (TPS) less than 50%, **without known actionable oncogenic targets***
 - Carboplatin or cisplatin, pemetrexed (Alimta), and pembrolizumab (Keytruda)†
 - Carboplatin or cisplatin, pemetrexed (Alimta), and cemiplimab-rwlc (Libtayo)
 - PD-L1 Expression (TPS) greater or equal to 50%, **without known actionable oncogenic targets***
 - Atezolizumab (Tecentriq)
 - Ineligible for Immunotherapy
 - Carboplatin, paclitaxel, and bevacizumab
 - Carboplatin or cisplatin and pemetrexed (Alimta)
- Squamous Cell Carcinoma Only
 - PD-L1 Expression (TPS) less than 50% **without known actionable oncogenic targets***
 - Pembrolizumab (Keytruda)†, carboplatin, and paclitaxel
 - Carboplatin, paclitaxel, and cemiplimab-rwlc (Libtayo)
 - Cisplatin, paclitaxel, and cemiplimab-rwlc (Libtayo)

Maintenance Therapy – Stages IIIB-IV, and Recurrent

- Non-Squamous Cell Carcinoma Only
 - Continuation bevacizumab
 - Continuation pemetrexed (Alimta)
 - Switch pemetrexed (Alimta)
 - If previously treated with carboplatin, pemetrexed, and pembrolizumab
 - Pembrolizumab (Keytruda)† and pemetrexed (Alimta)

* Actionable oncogenic targets refer to the driver aberrations in EGFR, ALK, and ROS1

† Administered at a dose of 200 mg every 3 weeks OR 400 mg every 6 weeks per the FDA label OR 2 mg/kg (up to a maximum of 200 mg) every 3 weeks, as clinically appropriate

Note: Pathways are independent of specific health plan medical policy coverage criteria. Health plan medical policy/clinical guidelines should be consulted to determine whether proposed services will be covered. Biosimilars of reference products listed are considered “on pathway.” However, reimbursement for biosimilar products may be impacted by health plan specific formularies, medical policy and preferred product rules.



8600 West Bryn Mawr Avenue
South Tower – Suite 800 Chicago, IL 60631

Last review: 11/5/2024 | Effective date: 1/20/2025

© 2025 Carelon Medical Benefits Management