Melanoma Pathways: Cutaneous Melanoma Patient Name: Date of Birth: Member Number: Treatment Start Date: Pathology:_ Stage:_ ICD-10 Code:____ Line of Therapy: **Biomarkers/Characteristics**: (Select all that apply) BRAF status: __V600E Mutation positive __V600K Mutation positive __Wild Type (no mutation) __Not Reported **Neoadjuvant Therapy** • Stages IIIB-IV (resectable) □ Pembrolizumab (Keytruda) □ Nivolumab (Opdivo) and ipilimumab (Yervoy) **Adjuvant Therapy** Stages IIB-III Resected ☐ Pembrolizumab (Keytruda) □ Nivolumab (Opdivo)

First Line of Therapy (1st Line)

- Stages IV and Recurrent
 - Any BRAF Status
 - □ Nivolumab (Opdivo)
 - ☐ Nivolumab (Opdivo) and ipilimumab (Yervoy)
 - □ Pembrolizumab (Keytruda)*
 - BRAF Mutated[†]
 - ☐ Encorafenib (Braftovi) and binimetinib (Mektovi)‡

- † BRAF mutations include V600E and V600K mutations
- ‡ First line only if the patient is not considered a suitable candidate for immunotherapy.

Note: Pathways are independent of specific health plan medical policy coverage criteria. Health plan medical policy/clinical guidelines should be consulted to determine whether proposed services will be covered. Biosimilars of reference products listed are considered "on pathway." However, reimbursement for biosimilar products may be impacted by health plan specific formularies, medical policy and preferred product rules.



^{*} Administered at a dose of 200 mg every 3 weeks OR 400 mg every 6 weeks per the FDA label OR 2 mg/kg (up to a maximum of 200 mg) every 3 weeks, as clinically appropriate